

# Traveling Electrical Waves in Cortex: Measurements with Optical Imaging Techniques and Insights from Phase Dynamics

David Kleinfeld, Department of Physics, UCSD, La Jolla, CA 92093-0319.

The theory of coupled phase oscillators [1] provides a framework to understand the emergent properties of networks of neuronal oscillators [2]. Such oscillators are a common feature in many brain areas [3]. When the architecture of the network is dominated by short-range synaptic connections, the pattern of electrical output is predicted to correspond to traveling plane and rotating waves of electrical activity, in addition to synchronized output. Here we discuss the experimental evidence for such patterns, and argue that the theory of coupled phase oscillators provides the foundation for understanding the traveling electrical waves that are observed in a variety of species.

We present the results of experiments on three preparations in which the issue of large-scale electrical activity was addressed, all of which involved the response to natural stimuli in the intact brain of awake animals. These include: (i) the central olfactory organ of the mollusk *Limax maximus* [4;5]; (ii) the primary visual cortex of turtle, *Pseudemys scripta* [6;7]; and (iii) the visuomotor cortices of cat [8]. The data collection involved the use of multiple-site electrical techniques, both with standard electrodes and with optical imaging technology in conjunction with voltage sensitive dyes [9].

The data reveal that waves are typically present during periods outside of stimulation, while synchronous or near-synchronous activity typically dominates in the presence of a strong stimulus [2]. We suggest that the continuum of phase-shifts during periods with traveling waves provide a means to scan and categorize the incoming sensory stream for novel features. Experiments to test the validity of the phase oscillator approach, as well as computational aspects of wave motion, are presented [2]. We further discuss possible mechanisms for the switch from patterned electrical activity to synchrony [10].

Supported by the NSF and the NIH/NCRR.

1. Kuramoto, Y., *Chemical Oscillations, Waves and Turbulence*. 1984, New York: Springer Verlag.
2. Ermentrout, G. B. and Kleinfeld, D., Traveling electrical waves in cortex: Insights from phase dynamics and speculation on a computational role. *Neuron*, 2001. **29**:1-12.
3. Basar, E. and Bullock, T. H., *Induced Rhythms in the Brain*. 1992, Boston: Birkhauser.
4. Delaney, K. R., Gelperin, A., Fee, M. S., Flores, J. A., Gervais, R., Tank, D. W., and Kleinfeld, D., Waves and stimulus-modulated dynamics in an oscillating olfactory network. *Proc. Natl. Acad. Sci. USA*, 1994. **91**:669-673.
5. Kleinfeld, D., Delaney, K. R., Fee, M. S., Flores, J. A., Tank, D. W., and Gelperin, A., Dynamics of propagating waves in the olfactory network of a terrestrial mollusk: An electrical and optical study. *J. Neurophysiol*, 1994. **72**:1402-1419.
6. Prechtl, J. C., Cohen, L. B., Mitra, P. P., Pesaran, B., and Kleinfeld, D., Visual stimuli induce waves of electrical activity in turtle cortex. *Proc. Natl. Acad. Sci. USA*, 1997. **94**:7621-7626.
7. Prechtl, J. C., Bullock, T. H., and Kleinfeld, D., Direct evidence for local oscillatory current sources and intracortical phase gradients in turtle visual cortex. *Proc. Natl. Acad. Sci. USA*, 2000. **97**:877-882.
8. Roelfsema, P. R., Engel, A. K., Konig, P., and Singer, W., Visuomotor integration is associated with zero time-lag synchronization among cortical areas. *Nature*, 1997. **385**:157-161.
9. Orbach, H. S., Cohen, L. B., and Grinvald, A., Optical mapping of electrical activity in rat somatosensory and visual cortex. *J. Neurosci.*, 1985. **7**:1886-1895.
10. Ermentrout, G. B., Flores, J., and Gelperin, A., Minimal model of oscillations and waves in the Limax olfactory lobe with tests of the model's predictive power. *J. Neurophysiol.*, 1998. **79**:2677-2689.